

INDIAN ASSOCIATION OF PHYSICS TEACHERS
NATIONAL STANDARD EXAMINATION IN CHEMISTRY 2018-19

Date of Examination: 25th November, 2018

Time: 11:00 to 13:00 hrs

Write the question paper code mentioned above on YOUR answer sheet (in the space provided), otherwise your answer sheet will NOT be assessed. Note that the same Q.P. Code appears on each page of the question paper.

Instructions to Candidates –

1. Use of mobile phones, smartphones, ipads during examination is **STRICTLY PROHIBITED**.
2. In addition to this question paper, you are given answer sheet along with Candidate's copy.
3. On the answer sheet, make all the entries carefully in the space provided **ONLY** in **BLOCK CAPITALS** as well as by properly darkening the appropriate bubbles.
Incomplete/incorrect/carelessly filled information may disqualify your candidature.
4. On the answer sheet, use **only BLUE or BLACK BALL POINT PEN** for making entries and filling the bubbles.
5. The email ID and date of birth entered in the answer sheet will be your login credentials for accessing performance report. Please take care while entering.
6. Question paper has 80 multiple choice questions. Each question has four alternatives, out of which **only one** is correct. Choose the correct alternative and fill the appropriate bubble, as shown.

Q. No. 22

a



c

d

7. A correct answer carries 3 marks whereas 1 mark will be deducted for each wrong answer.
8. Any rough work should be done only in the space provided.
9. Use of **non-programmable** calculator is allowed.
10. No candidate should leave the examination hall before the completion of the examination.
11. After submitting your answer paper, take away the Candidate's copy for your reference.

Please DO NOT make any mark other than filling the appropriate bubbles properly in the space provided on the answer sheet.

Answer sheets are evaluated using machine, hence CHANGE OF ENTRY IS NOT ALLOWED. Scratching or overwriting may result in a wrong score.

DO NOT WRITE ON THE BACK SIDE OF THE ANSWER SHEET.

Instructions to Candidates (continued) –

Read the following instructions after submitting the answer sheet.

12. Comments regarding this question paper, if any, can be shared only on Google forms, <https://goo.gl/forms/Lxb1l8Bqov3Cl9FQ2> till 27th November, 2018.

13. The answers/solutions to this question paper will be available on our website – www.iapt.org.in by 2nd December, 2018.

14. CERTIFICATES and AWARDS –

Following certificates are awarded by the IAPT to students successful in NSEs

(i) "Centre Top 10%" that will be sent to NSE centre by post.

(ii) "Statewise Top 1%" that can be downloaded after Feb -15th, 2019 from iapt.org.in

(iii) "National Top 1%". Certificates can be downloaded after Feb -15th, 2019 from iapt.org.in

15. Result sheets can be downloaded from our website in the month of February. The "Centre Top 10%" certificates will be dispatched to the Prof-in-charge of the centre by February, 2019.

16. List of students (with centre number and roll number only) having score above MAS will be displayed on our website (www.iapt.org.in) by 22nd December, 2018. See the Eligibility Clause in the Student's brochure on our website.

17. Students eligible for the INO Examination on the basis of selection criteria mentioned in Student's brochure will be informed accordingly.

18. Students qualified for OCSC (Chemistry) – 2019 will be awarded gold medals.

Useful Constants:

Charge of electron, $e = 1.602 \times 10^{-19}$ C

Mass of electron, $m_e = 9.1 \times 10^{-31}$ kg

Planck's constant, $h = 6.626 \times 10^{-34}$ J s


Speed of light, $c = 3.0 \times 10^8$ m s⁻¹

Avogadro constant, $N_A = 6.022 \times 10^{23}$ mol⁻¹


Molar gas constant, $R = 0.082$ L atm mol⁻¹ K⁻¹

$= 8.314$ J mol⁻¹ K⁻¹


1. Which of the energy values marked as I, II and III in the following diagram, will change by the addition of a suitable catalyst?





- (A) II only
(B) I and II
(C) II and III
(D) III only
2. The product 'X' in the following reaction is



- (A) a racemic mixture of ester
(B) an optically inactive ester
(C) an optically active ester
(D) a meso ester
3. At 298 K, change in internal energy for the complete combustion of fullerene, $C_{60}(s)$ an allotrope of carbon and the enthalpy of formation of $CO_2(g)$ are $-25970 \text{ kJ mol}^{-1}$ and -393 kJ mol^{-1} respectively. The enthalpy of formation of $C_{60}(s)$ at 298 K is
- (A) -2390 kJ
(B) $4.95 \times 10^4 \text{ kJ}$
(C) $2.60 \times 10^4 \text{ kJ}$
(D) 2390 kJ
4. Which of the following is not paramagnetic?
- (A) S^{2-}
(B) N^{2-}
(C) O^{2-}
(D) NO
5. Solubility product of AgCl is 1.8×10^{-10} . The minimum volume (in L) of water required to dissolve 1 mg of AgCl to close to
- (A) 0.5
(B) 7.5
(C) 50
(D) 0.75
6. The complex $[M(en)(Br)_2(Cl)_2]$ has two optical isomers. Their configurations can be represent as:






7. A sample of water from a river was analyzed for the presence of metal ions and the observations were recorded as given below:





Reagent added	Observation
dil HCl	No change
aq. Na ₂ CO ₃	White precipitate
aq. Na ₂ SO ₄	No change

The water sample is likely to contain


- (A) Ba²⁺ (B) Cu²⁺
 (C) Li⁺ (D) Mg²⁺
8. The lattice enthalpy and enthalpy of solution in water for solid NaCl are 753 kJ mol⁻¹ and 5 kJ mol⁻¹ respectively. If the solution enthalpies of Na⁺ and Cl⁻ are in the ratio 6 : 5, the enthalpy of hydration of Na⁺ ion is
 (A) 408 kJ mol⁻¹ (B) -412 kJ mol⁻¹
 (C) -408 kJ mol⁻¹ (D) -412 kJ mol⁻¹
9. The gaseous product obtained on reaction of BF₃ with LiH is
 (A) HF (B) H₂
 (C) B₂H₆ (D) F₂
10. The equilibrium constant K for the reversible reaction A = B is 2 × 10³ at 350 K. The rate constant of the forward reaction in the presence and absence of a suitable catalyst at the same temperature are 5 × 10⁴ s⁻¹ and 4 × 10⁻⁶ s⁻¹ respectively. The rate constant of the reverse reaction in the absence of the catalyst is
 (A) 4 × 10⁻³ s⁻¹ (B) 2.5 × 10⁻¹ s⁻¹
 (C) 1.6 × 10⁻⁷ s⁻¹ (D) 1.25 × 10⁻² s⁻¹
11. The number of stereoisomers possible for the following compound



- (A) 4 (B) 2
 (C) 16 (D) 32
12. An adsorption isotherm equation proposed by Langmuir is of the form $V = \frac{V_0 b P}{(1 + b P)}$, where V is the volume of gas adsorbed at pressure P. For a given adsorbate/adsorbent system, V₀ and b are constants. The dependent of V on P can be depicted as







13. For the reaction $4\text{NO}_2(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{N}_2\text{O}_5(\text{g})$, $\Delta H_{\text{reaction}} = -112\text{kJ}$. If the N_2O_5 is assumed to be formed in the reaction as a solid, $\Delta H_{\text{reaction}}$ will be ($\Delta H_{\text{sublimation}}$ of N_2O_5 is 54kJ mol^{-1})
- (A) -220kJ (B) -4kJ
 (C) -166kJ (D) -332kJ
14. Urea, $\text{CO}(\text{NH}_2)_2$ decomposes at 90°C as $\text{CO}(\text{NH}_2)_2(\text{aq}) \longrightarrow \text{NH}_4^+(\text{aq}) + \text{OCN}^-(\text{aq})$. Experimental data obtained for the reaction is given in the following plot.



- From the graph it can be inferred that
- (A) Average rate of the reaction is the same for successive time intervals of 10 h
 (B) unit of rate constant of the reaction is h^{-1}
 (C) rate constant of the reaction is the lowest at 30 h
 (D) the reaction is of zero order
15. If for an aqueous solution of a weak acid, $\text{pH} = \text{pK}_a + 2$ at 25°C , the approximate fraction of the acid in the dissociated form is
- (A) 1.1% (B) 0.99%
 (C) 99.0% (D) 9.9%

16. 2.0 L of N_2 gas kept at $25^\circ C$ and 5 atm pressure were expanded isothermally against a constant pressure of 1 atm until the pressure of the gas reaches 1 atm. Assuming ideal behaviour, reversible work of expansion in this process (in J) is close to
 (A) 810 J (B) -194 kJ
 (C) -810 kJ (D) 3390 kJ

17. The compound which would undergo a reaction with ammonia by S_N1 mechanism?




18. The daily energy requirement of a teenager is 7800 kJ. As calculated from the data given in the table below, the amount of glucose he has to consume (g) per day assuming that the entire energy he requires comes from the combustion of glucose is—

Molecule	ΔH_f (kJ mol $^{-1}$)
$C_6H_{12}O_6$	-1273
CO_2 (g)	-394
H_2O	-286

- (A) 262 (B) 500
 (C) 131 (D) 250
19. The pressure inside two gas cylinders of volume 25 m^3 and 50 m^3 are kPa and 20 kPa respectively. The cylinders are kept at the same temperature and separated by a valve. What is the pressure in the combined system when the valve is opened?
 (A) 30 kPa (B) 15 kPa
 (C) 16.7 kPa (D) 2.5 kPa
20. Aluminium and copper are extracted from their oxide and sulphide ores respectively. Which of the following is correct?
 (I) Copper is extracted by the auto reduction of copper oxide by copper sulphide
 (II) Aluminium cannot be obtained by chemical reduction due to its strong affinity for oxygen
 (III) In electrometallurgy of Al, graphite is used as cathode to avoid reoxidation of Al into Al_2O_3 by preventing formation of O_2
 (IV) Sulphide ores of copper are difficult to be reduced than the oxide ores
 (A) I, II and IV (B) II and III
 (C) II and III (D) II and IV

21. Which of the following graphs describes the relationship between $[H_3O^+]$ and $[OH^-]$ in an aqueous solution at a constant temperature?



22. From the given standard electrode potentials
 $Sn^{4+}(aq) + 2e^- \rightarrow Sn^{2+}(aq) \quad E^0 = 0.15 \text{ V}$
 $Br_2(l) + 2e^- \rightarrow 2Br^-(aq) \quad E^0 = 1.07 \text{ V}$
 The approximate free energy change for the process
 $2Br^-(aq) + Sn^{2+}(aq) \rightarrow Br_2(l) + Sn^{4+}(aq)$ is

- (A) 177.6 kJ (B) 355 kJ
 (C) -177.6 kJ (D) -355 kJ

23. Number of moles of $KClO_3$ that have to be heated to produce 1.0 L of $O_2(g)$ at STP can be expressed as

- (A) $1/3 (1/22.4)$ (B) $1/2 (1/22.4)$
 (C) $2/3 (1/22.4)$ (D) $3/2 (22.4)$

24. The sequence of reagents required for the following conversion is



- (A) (i) $B_2H_6/H_2O_2/OH^-$ (ii) Na (iii) C_2H_5I (B) (i) HCl (ii) C_2H_5ONa
 (C) (i) H_3O^+ (ii) Na (iii) C_2H_5OH (D) (i) H_3O^+ (ii) Na (iii) C_2H_5Cl



25. Among the following, number of oxygen atoms present is the maximum in

- (A) 1.0 g of O_2 molecules (B) 4.0 g of O atoms
 (C) 1.0 g of O_3 (D) 1.7 g of H_2O

26. Which of the following elements will exhibit photoelectric effect with light of the longest wavelength?

- (A) K (B) Rb
 (C) Mg (D) Ca

27. Compound 'X' in the following reaction is




28. The standard molar entropies of H_2 (g), N_2 (g) and NH_3 (g) are 130, 190 and 193 $\text{J mol}^{-1} \text{K}^{-1}$ respectively. For the reaction $\frac{1}{2}\text{N}_2(\text{g}) + \frac{3}{2}\text{H}_2(\text{g}) \rightleftharpoons \text{NH}_3(\text{g})$ ($\Delta H_{\text{reaction}} = -45 \text{ kJ}$) to be in equilibrium, the temperature must be equal to

- (A) 464 K (B) 928 K
(C) 737 K (D) 354 K

29. Density of CO_2 gas at 0°C and 2.00 atm pressure can be expressed as

- (A) 2 g m^{-3} (B) 2 g m^{-3}
(C) $4 \times 10^3 \text{ kg m}^{-3}$ (D) 8 g L^{-1}

30. The maximum number of moles of CH_3I consumed by one mole of crixivan, a drug used against AIDS is



Crixivan





- (A) 2 (B) 3
(C) 5 (D) 7

31. Concentration of K^+ ions inside a biological cell was found to be 25 times higher than that outside. The magnitude of the potential difference between the two sides of the cell is close to (2.303 RT/F) can be taken as 59 mV; difference in concentrations of other ions can be taken as negligible).

- (A) 4.2 mV
(C) 82 mV
- (B) 195 mV
(D) -82 mV

32. The standard redox potential for the reaction $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$ is -1.23V . If the same reaction is carried out at 25°C and at $\text{pH} = 7$, the potential will be
(A) -0.82V
(B) -3.28V
(C) 0.82V
(D) -1.18V


33. The order of pK_a values of the following acids is



- (A) $\text{IV} > \text{I} > \text{III} > \text{II}$
(B) $\text{III} > \text{IV} > \text{I} > \text{II}$
(C) $\text{II} > \text{I} > \text{III} > \text{IV}$
(D) $\text{II} > \text{III} > \text{I} > \text{IV}$

34. If the radius of the hydrogen atom is 53 pm , the radius of the He^+ ion is close to
(A) 75 pm
(B) 38 pm
(C) 106 pm
(D) 27 pm

35. A substance X was heated at constant pressure and the temperature observed at various times of heating was plotted as given below:








Which of the following is/are correct?

- I. Melting point X is -5°C
 II. Solid and liquid forms of X coexist in the region b
 III. Boiling point of X is 55°C
 IV. Solid and liquid forms of X coexist in the region d

(A) I and IV
(C) III only

(B) II and III
(D) I, II and III

36. The major product of the following reaction is



37. In which of the following, all the bond lengths are not the same?

- I. IF₄⁺ II. BF₄⁻ III. SF₄ IV. TeCl₄
(A) I, II, IV (B) II, III, IV
(C) I, III, IV (D) I, II, III


38. Among the following, the reaction/s that can be classified as oxidation – reduction is/are

- I. Cr₂O₇²⁻ (aq) + 2 OH⁻ (aq) → 2 CrO₄²⁻ + H₂O(l)
II. SiCl₄ (l) + 2Mg(s) → 2MgCl₂ (l) + Si(s)
III. 6Cl₂ (l) + 12KOH(l) → 2KClO₃ (s) + 10KCl + 6H₂O(l)
IV. 2H₂O₂ → 2H₂O(l) + O₂ (g)
(A) I and IV (B) I, II and III
(C) II, III and IV (D) IV only

39. Among the following pairs, the one in which both the compounds as pure liquids can show significant auto ionization is

- (A) H₂O and H₂S (B) BrF₃ and ICl₃
(C) PF₅ and PCl₅ (D) HF and HCl


40. The number of quaternary and chiral carbon atoms present in elatol, isolated from an alga are respectively



Elatol

- (A) 2, 3 (B) 4, 2
(C) 3, 2 (D) 1, 3

41. Compound X ($pK_a \sim 15$) and Y ($pK_a \sim 10$), both produce H_2 on treatment with sodium metal and both yield a mixture of isomers on mononitration. X and Y respectively are:




- (A) IV, I
(B) III, II
(C) III, I
(D) I, III

42. A crystal of KCl containing a small amount of $CaCl_2$ will have
(A) vacant Cl^- sites
(B) vacant K^+ sites and a higher density as compared to pure KCl
(C) vacant K^+ sites and a lower density as compared to pure KCl
(D) K^+ ions in the interstitial sites

43. In the following reaction, the values of a, b and c, respectively ly
 $aF_2(g) + bOH^-(aq) \rightarrow cF^-(aq) + dOF_2(g) + eH_2O(l)$
 (A) 3, 2, 4
(B) 3, 4, 2
(C) 2, 2, 4
(D) 2, 2, 2


44. The monosaccharide present in the following disaccharide is



- (A)
- (B)
- (C)
- (D)

45. The IUPAC name of the complex $[\text{Pt}(\text{en})(\text{NH}_3)(\text{Cl})_2(\text{ONO})][\text{Ag}(\text{CN})_2]$ is
 (A) monoamminedichlorido (ethane -1, 2, diammine) nitritoplatinum (IV) dicyanoargentate (I)
 (B) monoaminebischlorido (ethane -1, 2 - diammine) nitroplatinate (IV) dicyanosilver (I)
 (C) monoaminebischlorido (ethane -1, 2 - diammine) nitritoplatinate (IV) dicyanoargentate (I)
 (D) monoamminedichlorido (ethane -1, 2 - diammine) nitritoplatinum (IV) dicyanoargentate (I)

46. The correct order of basicity of the following species is



- (A) III < IV < II < I
 (B) III < I < II < IV
 (C) III < II < I < IV
 (D) IV < I < II < III


47. Which among the following is nonlinear?

- (A) N_3^-
 (B) ClF_2^-
 (C) Br_3^-
 (D) BrCl_2^+

48. The compound most likely to lose water on protonation is



49. The Newman projection shown is the same as




- (A) I and IV
 (B) II and III
 (C) III and IV
 (D) I and II

50. Which one of the following is not used as a monomer for the synthesis of a high molecular weight silicone polymer?


- (A) MeSiCl_3
 (B) Me_2SiCl_2
 (C) Me_3SiCl
 (D) PbSi_3

51. In $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$, a superconducting oxide that got George Bednorz and Karl Mullar the Nobel prize in 1986. Cu can exist in both +2 and +3 oxidation states and their proportion depends on the value of 'x'. In $\text{YBa}_2\text{Cu}_3\text{O}_{7.0.5}$
- (A) 0.5 moles of Cu are in +3 oxidation state
 (B) 5% of Cu is in +3 oxidation state
 (C) All the Cu is in +3 oxidation state
 (D) All Cu is in +2 oxidation state


52. Compound 'Y' (molar mass = 88.1 g mol^{-1}) containing 54.52% carbon, 9.17% hydrogen and 36.31% oxygen gives a reddish – brown precipitate in Fehling's test. 'Y' is



53. The IUPAC name of the following compound is




- (A) 1 – Bromo – 4 – chloro – 3 etheny ibutane
 (B) 4 – Bromon – 1 – chloro – 3 ethenylbutane
 (C) 3 – (Bromomethyl) – 5 – chloropent – 1 – ene
 (D) 3 – (Bromomethyl) – 1 – chloropent – 4 – ene
54. The correct order of boiling points of the following compounds is




- (A) III < IV < II < I
 (B) I < III < IV < II
 (C) I < II < III < IV
 (D) IV < III < I < II
55. Which of the following is a strong oxidizing agent?
- (A) AlCl_3
 (B) TiCl_3
 (C) NF_3
 (D) PCl_3

56. The molecule in which all atoms are not coplanar is




57. The most stable radical among the following is


(A)




(B)



(C)



(D)



58. During World War II, soldiers posted at high altitudes experienced crumbling of the tin buttons of their uniforms into a grey powder. This can be attributed to


(A) oxidation of tin

(B) interaction with nitrogen in the air at low pressure


(C) change in the crystal structure of tin

(D) reaction of tin with water vapour in the air


59. The molecules that can exhibit tautomerism are




I



II



III



IV

(A) I, IV
(C) III, IV

(B) II, III
(D) I, II

60. A scientist attempts to replace a few carbon atoms in 1.0 g of diamond with boron atoms or nitrogen atoms in separate experiments. Which of the following is correct?


(A) The resulting material with B doping will be an n type semiconductor


(B) The resulting material with B doping will be a p type semiconductor

(C) B doping is NOT possible as B cannot form multiple bonds




(D) The resulting material with N doping will be a p – type semiconductor

61. Compound 'P' that undergoes the sequence of reactions given below to give the product Q is











62. The most stable Lewis structure of N_2O is





63. The major product 'X' formed in the following reaction is



64. Which of the following accounts best for the fact the F^- is smaller than O^{2-} ?



65. The correct sequence of reagents from those listed below for the following conversion is—





- (A) IV – I – III (B) III – IV – I
 (C) II – I – III (D) I – II – III


66. An orbital among the following that has two radial nodes and two angular nodes is

- (A) 3d (B) 4p
 (C) 4f (D) 5d


67. The compound 'X' undergoing the following reaction is




(A)



(B)



(C)





(D)

68. A dilute solution of an alkali metal in liquid ammonia is

- (I) blue in color (II) conducts electricity
 (III) paramagnetic (IV) an oxidising agent

- (A) I and III (B) II and IV
 (C) I and III (D) I and III

69.



- (A) i, ii and iii (B) i and ii
 (C) i and iii (D) ii and iii


70. The C–O bond length is the shortest in

- (A) $[\text{Cr}(\text{CO})_6]$ (B) $[\text{Mo}(\text{CO})_6]$
 (C) $[\text{Mn}(\text{CO})_6]^+$ (D) $[\text{V}(\text{CO})_6]^-$

71. The rate of the reaction between two reactants X and Y can be expressed as $R = k[X]^2[Y]$. In an experiment, the initial rate of the reaction was found to be R_1 when the initial concentration of X and Y are $[X_0]$ and $[Y_0]$. Another experiment was performed in which $[X_0]$ was taken as $\frac{1}{2}[X_0]$. What should be $[Y_0]$ in this experiment to get the initial rate as $0.5R_1$?

- (A) $4[Y_0]$ (B) $\frac{1}{2}[Y_0]$
 (C) $2[Y_0]$ (D) $[Y_0]$

72. Among the following, the compound that has the highest dipole moment is
 (A) $\text{CH}_3\text{COOCH}_3$ (B) CH_3CONH_2
 (C) $\text{CH}_3\text{COC}_2\text{H}_5$ (D) CH_3COCl
73. A common method to clean acid spills is to use Na_2CO_3 (Molar mass 106g). If 50.0 mL of 0.75 M HCl is split on a wooden surface, the amount of Na_2CO_3 required is
 (A) 3.75 g (B) 7.5 g
 (C) 2.0 g (D) 4.0 g
74. The spin-only magnetic moments of $[\text{Fe}(\text{NH}_3)_6]^{3+}$ and $[\text{FeF}_6]^{3-}$ (in units of BM) respectively are
 (A) 1.73 and 1.73 (B) 5.92 and 1.73
 (C) 1.73 and 5.92 (D) 5.92 and 5.92
75. The major product of the following reaction is



76. The standard electrode potential (E^0) of the Daniel cell is 1.1 V and the overall cell reaction can be represented as $\text{Zn}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu}(\text{s})$.
 (A) 1.0 M Zn^{2+} , 1.0M Cu^{2+} (B) 1.2 M Zn^{2+} , 1.2 M Cu^{2+}
 (C) 0.1 M Zn^{2+} , 1.0M Cu^{2+} (D) 1.0 M Zn^{2+} , 0.01M Cu^{2+}
77. Penicillamine is used in the treatment of arthritis. One molecule of penicillamine contains a single sulphur atom and the weight percentage of sulphur in penicillamine is 21.49%. Molecular weight of penicillamine in g mol^{-1} is
 (A) 85.40 (B) 68.76
 (C) 125.2 (D) 149.2
78. An ion exchange resin, RH_2 can replace Ca^{2+} in hard water as $\text{RH}_2 + \text{Ca}^{2+} \rightarrow \text{RCa}^{2+} + 2\text{H}^+$
 When a 1.0 L hard water sample was passed through the resin, all H^+ ions were replaced by Ca^{2+} ions and the pH of eluted water was found to be 2.0. The hardness of water (as ppm of Ca^{2+} in the sample of water treated is
 (A) 50 (B) 100
 (C) 125 (D) 200
79. The analysis of three different binary oxides of bromine (Br) and oxygen (O) gives the following results:

FREE